

Series paper #4

Economics of growing slash and loblolly pine on a 33-year rotation with and without thinning, fertilization, and pine straw—net revenue and rate of return

August 2007

E. David Dickens, Coleman W. Dangerfield Jr., and David J. Moorhead; Forest Productivity Associate Professor, Forest Economics Professor-Emeritus, and Silviculture Professor, respectively, with the Daniel B. Warnell School of Forest Resources The University of Georgia, Athens, Georgia 30602-2152

Abstract

Since early 1998 forest industry, forestland ownership, global markets, and wood supply and demand (pulpwood, sawtimber, chips, etc.) regionally and world-wide have changed dramatically. Non-industrial private forest (NIPF) landowners have realized reduced product market availability and increased price uncertainty during this period in the southeastern United States. Lower Atlantic and Gulf Coastal Plain NIPF landowners seek management options utilizing two commonly available pine species; loblolly (*Pinus taeda* L.) and slash (*Pinus elliotii*, Engelm.) to enhance feasibility, profitability, and cash-flow of production forestry enterprises. At the same time, NIPF landowners desire heightened flexibility across time required to achieve marketable forest products. This paper examines feasibility, profitability, and cash-flow of mixed product class rotation management options of 33-year rotation scenarios affecting wood-flow for slash and loblolly pine plantations including thinning, fertilization, and pine straw harvests under alternative levels of productivity, establishment costs, and product prices. Financial measures of profitability calculated include net revenue and rate of return (ROR). Soil expectation value (SEV) and annual equivalent value (AEV) are discussed in series paper #5 for the 33-year rotation scenarios.

Introduction

Private non-industrial forest (NIPF) landowners in the Atlantic and Gulf Coastal Plain from South Carolina to Mississippi question whether to plant slash or loblolly pine on cut-over and old-field sites. They also question spending moderate to relatively large sums of money in intensive forest management under the current and anticipated stumpage prices and economic uncertainty. Many NIPF landowners speculate how attractive chip-n-saw / sawtimber rotations with thinning are for southern pines. To address these questions, we used the Georgia Pine Plantation (GaPPs 4.20) growth and yield Model developed by Bailey and Zhao (1998).

Methodology

Common assumptions

The rotation age was set at 33-years for slash and loblolly pine plantations. Net revenue was calculated as the sum of all revenues minus all costs in today's dollars. Rate of return (ROR) was calculated after all intermediate costs (fertilization) and

returns (pine straw) were discounted at 8 percent. Fire protection cost was assumed \$2/ac/yr, stand management at \$2/ac/yr, and property taxes at \$5/ac/yr. Thus, the total annual costs for each year of the rotation were \$9/acre. This value cost goes in the transaction table as an annual cost during the rotation. The present value of this net, annual cost flow is \$103.63 during the 33-year rotation. Results are reported in constant dollars, before taxes. It is assumed that land is already owned.

Site Preparation and Planting Costs

Three site preparation and planting (SP+PL) costs rise in increments of \$125/acre (\$125, \$250 and \$375/acre). These costs represent the following site preparation and planting scenarios:

- ▶ The relatively low site preparation and planting cost of \$125/acre could include machine planting and the use of a post plant herbicide to control herbaceous weeds on an old-field site or glyphosate @ 1 gallon/ac or prescribe burning (low level) site preparation and roughland planting on a cutover site.
- ▶ The more moderate (\$250/acre) establishment cost could include a mechanical site prep treatment, burn and plant or a herbicide, burn, plant, and herbaceous weed control (Dubois and others. 1999).
- ▶ The higher (\$375/acre) establishment cost could include a combination of chemical and mechanical site preparation as can be the case on many flatwoods cutover sites.

Site preparation options and associated costs vary extensively by location, prior stand history, harvesting utilization, landowner objectives, monies available, and anticipated future stumpage value and demand. The assumption used was that level of site preparation intensity was matched to level of competition control needed so that wood-flows were comparable within site productivity levels, after site preparation and planting.

Product class specifications

Product class specifications are:

- ▶ pulpwood (PW) at a d.b.h. of 4.6 to 9 inches to a 3 inch top;
- ▶ chip-n-saw (CNS) at a d.b.h of 9 through 12 inches to 6 inch top; and,
- ▶ sawtimber (ST) with a d.b.h greater than 12 inches to a 10 inch top (inside bark) were assumed (Table 1).

Georgia stumpage Prices, reported through Timber Mart-South[©] (TM-S) for 1st quarter year 2004 average, used in this analysis for loblolly and slash were net of property taxes at harvest (2.5 percent) and net of marketing costs (8 percent). The low TM-S prices for pulpwood and chip&saw were used for thinning prices and average TM-S prices for pulpwood, CNS, and ST are used for the clearcut. Cash and net converted prices are found in Table 2. It is assumed that land is owned throughout the scenarios.

Species specific assumptions

The slash pine scenarios assumed 500 living trees per acre (TPA) at age 5-years-old. A base mean annual increment of 1.91 cd/ac/yr (5.26 tons/ac/yr) @ age 33-years-old without fertilization and thinning was assumed (Table 7). The base slash scenario woodflow was 12 percent less than base loblolly woodflow (Shiver and others 1999) at age 33-years. The assumed fertilizer applications increased merchantable volume by an average of 0.50 cd/ac/yr (1.38 tons/ac/yr) for eight to ten years following treatment (Jokela and Stearns-Smith 1996).

The loblolly pine survival was assumed to be 500 TPA at age 5-years-old. The base mean annual increment for loblolly was assumed to be 2.15 cds/ac/yr (5.95 tons/ac/yr, Table 8) through age 33-years-old without fertilization or thinning. The base loblolly woodflow was 12 percent greater than the slash base woodflow (Shiver and others 2000) at age 24-years. The assumed fertilizer applications aggressively increased merchantable volume by an average of 0.65 cd/ac/yr (1.79 tons/ac/yr) for eight to ten years (NCSUFNC 1998). Figures 1, 2, and 3 show wood flow for scenarios #1, 2, and 3 (scenario #4, 5, and 6 have same wood flows as #1, 2, and 3, respectively).

Scenarios for the 33-year Rotation

The following are the seven slash (Table 7) and loblolly (Table 8) pine scenarios:

- (1) thin @ age 15- and 24years to 65 ft²/ac, no straw, no fertilization ,
- (2) thin @ age 15- and 24years to 65 ft²/ac, no straw, fertilize @ age 25-years,
- (3) thin @ age 15- and 24years to 65 ft²/ac, no straw, fertilize @ ages 16- and 25-years,
- (4) thin @ age 15- and 24 years to 65 ft²/ac, pine straw @ \$50/ac/yr from age 8-through age 14-years, no fertilization,
- (5) thin @ age 15- and 24 years to 65 ft²/ac, pine straw @ \$100/ac/yr from age 8-through age 14-years, fertilize @ age 6-years,
- (6) thin @ age 15- and 24 years to 65 ft²/ac, pine straw @ \$100/ac/yr from age 8-through age 14-years, fertilize @ age 6- and 25-years, and
- (7) thin @ age 15- and 24 years to 65 ft²/ac, pine straw @ \$100/ac/yr from age 8-through age 14-years, clean-up in year 16, pine straw @ \$50/ac/yr from age 17-through age 23-years, fertilize @ age 6-, 16-, and 25-years.

Forest management activities

Thinning

All scenarios for both pine species include two thinnings; @ age 15- and 24-years-old. Residual basal area (RBA), after thinning (5th row with selection from below) is set at 65 sq. ft/ac.

Fertilization

Either no fertilization (scenario #1 and 4) or a fertilizer and application cost of \$100/ac for slash and loblolly per application at age 6-, 16-, and/or 25-years were assumed. Fertilization with 150 then 200 N + 40 P (as diammonium phosphate and urea) per acre was part of this scenario to maintain pine straw production rates (Dickens 1999), to enhance wood volume (NCSUFNC 1998), and change product class distribution (Peinaar and Rheney 1996, Dickens 2001). Fertilization timing at age 6-years-old was two years prior to the initiation of straw raking (just prior to canopy closure). The second application, ten years later, was just after a thinning (thinning scenario) and after the response (wood and straw) to the first application has become negligible. The 25-year fertilizer application was made to increase sawtimber volume. The periodic fertilizer application costs are converted to present values (PV) in year one, then re-computed as annual equivalent values (AEV). These AEVs were then put in the transaction table as annual expense cash-flows (Table 3).

Scenarios with fertilization for both loblolly and slash pine were set-up as follows: (#2) to delay fertilization cost and/or to promote additional growth on best (leave) trees only after the 2nd thinning (age 25-years only), (#3) to fertilize residual trees after each thinning (ages 16- and 25- years), (#4 & 5) to maintain or enhance pine straw production from canopy closure (age 6-years only), (#6) to maintain pine straw production (age 6- and 25-years) to the 1st thinning then put more growth on sawtimber trees after 2nd thin, and (#7) to maintain or enhance pine straw production from just prior to canopy closure (age 6-yr) to the first thinning and to change product class distribution and put extra growth on best trees after 1st and 2nd thinning age 16- and 25-years, (Table 7 and 8).

Pine straw

The pine straw income assumptions included were as follows: \$50 and \$100/ac/yr raking income for the slash and loblolly scenarios has been noted in south (slash) and central (loblolly) Georgia between 1998 and 2003 (Doherty 2004). Pine straw is raked starting in year 8 (approximating canopy closure) for slash and loblolly pine. Periodic pine straw income was converted to present values (PV) in year one, then re-computed as annual equivalent values (AEV) at the discount rate of 8 percent. These AEVs were then put in the transaction table as annual income cash-flows (Table 4).

Typically pine straw raking in Georgia ceases after the first thinning due to large understory vegetation growth in thinned stands and the abundance of unthinned, relatively clean loblolly and slash pine stands available. Yet many acres of thinned loblolly and longleaf stands in South and North Carolina are raked. Some pine straw contractors in Georgia anticipate that some thinned loblolly, longleaf, and slash pine stands may be rakeable in the future (supply and demand). Therefore, we included a scenario for loblolly and slash pine with raking re-commencing two years after thinning at half the income rate prior to the thinning. There was an associated clean-up cost to get the stand rakeable of \$70/acre (Table 5). Scenarios that included pine straw income for both species are #s 4 through 7 (Table 7 and 8).

Results

When considering results, more weight should be given to the relative differences among scenarios than to the numeric values calculated for any one scenario. Among-scenario differences are more important than absolute values because scenarios were analyzed under a common set of conservative assumptions. Values realized for your individual case may be greater or less than those of this study. However, the relative magnitude of difference between and ranking of the scenarios should remain the same under a different set of underlying assumptions.

Net revenues and rates of return ranges

Net revenues (NR) ranged from \$2,850 (base slash pine scenario with highest site prep and plant cost) to \$5,581 per acre (loblolly with pine straw @ \$100/ac/yr; age 8-14 yrs and \$50/ac/yr age 17-23 yrs, fertilize three times with lowest site prep and plant cost (Table 7 and 8). Ranking of scenarios by NR within a SP+PL level were as follows: 7 > 6 > 5 > 4 ≈ 3 > 2 > 1 for slash pine. Ranking of scenarios by NR within a SP+PL level were as follows: 7 > 6 > 5 > 3 > 4 > 2 > 1 for loblolly pine. Net revenues for slash pine (growing at approximately 15 percent less than loblolly) were 12 to 17 percent less than corresponding loblolly scenarios except for scenario #1 (Table 7 and 8).

Rates of return (ROR) for all scenarios (42 scenarios in all) for both species ranged from 6.69 and 7.18 percent (base slash and loblolly scenarios with highest site preparation and planting cost, respectively) to 16.89 and 17.31 percent (slash and loblolly pine scenarios with the lowest site preparation and planting cost, fertilize three times, and rake straw @ \$100/ac/yr from age 8- through 14-years and \$50/ac/yr from age 17- through age 23-years, respectively) using the aforementioned assumptions (Table 7 and 8). Ranking of scenarios by ROR (within a SP+PL level) were as follows: 7 > 6 > 5 > 4 > 3 > 2 > 1 for loblolly pine (with the 0.75 cd/ac/yr fertilizer response). Ranking of the slash pine scenarios by ROR (within a SP+PL level) were as follows: 7 > 6 > 5 > 4 > 3 ≈ 2 > 1 (with the 0.40 cd/ac/yr fertilizer response).

Impact of pine straw income on net revenues and rates of return

Pine straw income prior to thinning (age 8-14-yrs) increased NR by \$350, (scenario #4, pine straw only), \$867 (scenario #5, straw + extra wood), and \$1065/acre (scenario #6, straw + extra wood) compared to the no pine straw, no fertilization scenario (#1) for slash pine (Table 7). Pine straw income prior to thinning (age 8-14-yrs) increased NR by \$350, (scenario #4, pine straw only), \$1057 (scenario #5, straw + extra wood), and \$1408/acre (scenario #6, straw + extra wood) compared to the no pine straw, no fertilization scenario (#1) for loblolly pine (Table 8). When pine straw was performed before and after the thinning, (scenario #7) NR increased by \$1636 and \$2052/acre (\$750/acre in net straw minus fertilization income) over the no pine straw, no fertilization base scenario (#1) for slash and loblolly, respectively (Table 7 and 8).

The addition of pine straw income (@ \$50/ac/yr from age 8 – 14 years) for slash pine in scenario #4 increased base scenario (#1) rates of return from 6.69, 7.91, and 9.86

percent to 8.08, 9.75, and 12.88 percent for the \$375, \$250, and 125/ac SP+PL costs, respectively (Table 7). Raising the annual pine straw income for slash pine to \$100/ac/yr from age 8- through 14-years and fertilizing twice (scenario #6) increased rates of return to 9.34, 11.30, and 15.32 percent for the @ \$375, \$250, and \$125/ac SP+PL costs, respectively (Table 7). Pine straw raking in the slash scenario prior to thinning (age 8 through 14-years) at \$100/ac/yr and after the 1st thinning (ages 17- through 23-years) at \$50/ac/yr (scenario #7) produced rates of return of 10.06, 12.22, and 16.89 percent for the \$375, \$250, and \$125/ac SP+PL costs, respectively (Table 7).

The addition of pine straw income (@ \$50/ac/yr from age 8 – 14 years) for loblolly pine in scenario #4 increased base scenario (#1) rates of return from 7.18, 8.42, and 10.41 percent to 8.52, 10.20, and 13.35 percent for the \$375, \$250, and 125/ac SP+PL costs, respectively (Table 8). Raising the annual pine straw income for loblolly pine to \$100/ac/yr from age 8- through 14-years and fertilizing twice (scenario #6) increased rates of return to 10.10, 12.12, and 16.26 percent for the \$375, \$250, and \$125/ac SP+PL costs, respectively (Table 8). Pine straw raking in the loblolly scenario prior to thinning (age 8 through 14-years) at \$100/ac/yr and after the 1st thinning (ages 17- through 23-years) at \$50/ac/yr (scenario #7) produced rates of return of 10.67, 12.81, and 17.31 percent for the \$375, \$250, and \$125/ac SP+PL costs, respectively (Table 8).

Impact of fertilization on net revenues and rates of return

Net revenues increased by \$167 and \$357/acre with fertilization at age 25-years (scenario #2) compared to the no fertilization scenario (#1) for slash and loblolly pine, respectively (Table 7 and 8). Fertilization just after both thinnings (scenario #3) increased NR by \$365 and \$708/acre compared to the thin only scenario (#1) (Table 7 and 8) for loblolly and slash pine, respectively.

Fertilization in slash pine stands with 200 N + 40 P/acre after the first thinning increased ROR by 0.20 to 0.23 percent across the three SP+PL levels and loblolly ROR by 0.07 to 0.20 percent (scenario #1 vs #2, Table 7 and 8). Fertilization at age 16-and 25-years ROR (scenario #3) was 0.05 to 0.27 percent above the ROR for the thin only scenario (#1) for slash pine and about 0.53 to 0.72 percent above the thin only scenario for loblolly pine.

Impact of establishment costs on net revenues and rates of return

The impact of establishment costs (site preparation and planting; SP+PL) was straightforward; net revenues differing by increments of \$125/acre within a scenario by species. Establishment cost impact on the time-value of money though was large.

The impact of establishment costs within a management level (scenario) was large enough to illustrate the importance of choosing the right SP+PL for a given site. The impact of SP+PL on rates of return became larger as management inputs increased for both species. For example: the base slash pine scenario (#1) of thin twice, no fertilization, no pine straw had ROR values of 6.69, 7.91, and 9.86 percent for the \$375, \$250 and \$125/acre SP+PL costs, respectively with differences of 1.22 and 3.17

percent. Slash pine scenario #7 had ROR values of 10.06, 12.22, and 16.89 percent, differences of 2.16 and 6.83 percent (Table 7). The impact of SP+PL in the loblolly scenarios showed the same trend as the slash pine scenarios. The base loblolly pine scenario (#1) ROR values were 7.18, 8.42, and 10.41 percent for the \$375, \$250 and \$125/acre SP+PL costs, respectively with differences of 1.24 and 3.23 percent (Table 8). Loblolly pine scenario #7 ROR values were 10.67, 12.81, and 17.31 percent; differences of 2.14 and 6.64 percent.

Impact of management inputs on net revenues and rates of return

Generally, increasing management, whether through fertilization or with pine straw increased net revenue and rates of return for both species. Fertilization improved net revenue by \$167 and \$365/acre for slash pine and by \$357 and \$708/acre for loblolly pine (scenario #2 and 3 vs #1, Table 7 and 8). Adding pine straw increased NR by a wide margin; \$350 to \$1,636/acre for slash pine and by \$350 to \$2,052/acre (pine straw + extra wood) for loblolly pine (scenario #4 and 7 vs #1, Table 7 and 8).

Fertilization after the 2nd thinning (age 25-years, scenario #2) improved ROR for both species by a negligible amount (0.06) to 0.23 percent over the unfertilized scenario (#1, Table 7 and 8). Fertilization after the 1st and 2nd thinning (scenario #3) increased ROR by a small amount (0.05 percent) to 0.72 percent over the unfertilized scenario (#1) for both species. Even though fertilization did not increase ROR greatly, remember NR increased significantly.

The \$50/ac/yr pinestraw income from age 8- through 14-years greatly improved ROR for both species; by 1.39 to 3.02 percent for slash pine (scenario # 1 vs 4) and 1.34 to 2.94 percent for loblolly (scenario #1 vs 4) (no fertilization, Table 7 and 8). The \$100/ac/yr pine straw revenue from age 8- through age 14-years further improved ROR by 1.18 to 2.64 percent for slash pine (scenario #4 vs 5) and by 1.07 to 2.21 percent for loblolly pine (scenario #4 vs 5) over the \$50/ac/yr income rate (Table 7 and 8).

Summary

Wood flow, fertilization responses, and pine straw

The 1.91 (5.26 tons/ac/yr) and 2.15 cd/ac/yr (5.95 tons/ac/yr) base productivity levels through age 33-years-old for slash and loblolly, respectively, are very realistic on most cut-over sites with chemical site preparation and post-plant herbaceous weed control (Pienaar and Rheney 1996) and may be conservative on most old-field sites.

Exceptions would be problem soils such as deep sands (Typic Quartzipsammments) of the Sand Hills or shallow, rocky soils of the Piedmont physiographic region.

These scenarios do illustrate that if the aforementioned base growth rates for slash pine and loblolly pine are assumed then the establishment expenditures (site preparation and planting costs) need to be used wisely. In many cases the establishment phase decisions (site preparation type, timing, and quality, site preparation effects on near- or long-term site productivity, woody and herbaceous weed

control efficacy, species selection, seedling genetics and size, seedling survival) can improve growth rates above those used here, therefore improving rates of return.

The fertilization response for slash pine of an average of 0.50 cd/ac/yr (1.38 tons/ac/yr) for 8 to 10 years (Jokela and Stearns-Smith 1993, Martin and others 1999) is realistic assuming the site is deficient in N or N+P and the genetics and stocking are not response limiting. The fertilization response average of 0.65 cd/ac/yr (1.79 tons/ac/yr) for loblolly pine over 8 to 10 years is slightly above an average response (NCSFNC 1999), therefore the site must be very deficient in N or N+P, leaf area is low and genetic and stocking are not growth limiting. No increase in pine straw income per acre was assumed with fertilization. Fertilization studies (Blevins and others 1996, Dickens 1999) illustrate that pine straw production can be increased by an average of 40 to 50 percent over unfertilized stands on marginal fertility soils. Fertilization was included in the pine straw production scenarios to maintain straw production as nutrients are removed/displaced with each raking. Adding potassium (K), magnesium (Mg), sulfur (S), calcium (Ca) and/or micro-nutrients (boron, manganese, or copper) to the above assumed N+P application(s) may be necessary based on site and stand needs which should increase fertilization cost.

When wood value only is considered, loblolly produced more wood, more wood value (a diameter driven function), a higher NR and ROR is realized with the aforementioned assumptions. Recent studies (Shiver and others 1999) have shown that loblolly will grow more wood than slash on a number of soils where both species are grown. Loblolly's superior wood volume yields do not necessarily equate to higher per acre or per unit wood stumpage prices. Clark (2002) noted that slash pine yielded more number one lumber, had a slightly greater (4 to 11 percent greater) density, and 4 percent less moisture content than loblolly pine in growing in the same stand.

Discussion

Non-industrial private forest landowners do have some attractive forest management options with both slash and loblolly pine even when using low to medium stumpage prices. Generally, increasing forest management activities (thinning, fertilization, adding pine straw) increased rates of return at the wood growth increments used.

If a rate of return of 8 percent or better is a landowner goal with the stumpage prices (Georgia 1st Qtr 2004, TM-S 2004) and the wood production rates used, then all scenarios for loblolly and scenarios #2 through 7 for slash pine at the lower two site preparation and planting establishment costs achieved that. At the highest SP+PL level, an ROR of 8 percent or better was achieved only when pine straw income @ \$50/ac/yr or better was realized for both species (scenarios #4 through 7, Table 7 and 8).

If a rate of return of 10 or better is a landowner objective under the aforementioned assumptions, then pine straw production to achieve \$100/ac/yr (scenario #6 and 7) for loblolly and \$100/ac/yr then \$50/ac/yr (scenario #7) for slash pine is required @ the

highest SP+PL level cost. A 10 percent or better ROR can be realized at the moderate SP+PL cost with the rake @ \$100/ac/yr scenarios (#5 - 7) for slash pine (Table 7) and @ the \$50 or \$100/ac/yr rake income for loblolly pine (scenario #4-7, table 8). At the lowest SP+PL cost all scenarios but scenario #1 and 3 for slash pine had an ROR greater than 10 percent.

A rate of return of 12 percent or better was not achieved for either species at the highest SP+PL level in any of the scenarios. A rate of return of 12 percent or better was realized with the \$100/ac/yr pine straw income (loblolly) or the \$100 then \$50/ac/yr pine straw income (slash and loblolly pine) at the moderate SP+PL costs (scenario #6 and 7, Table 7 and 8). At the low SP+PL cost, an IRR of 12 percent or better was realized with all the pine straw income @ \$50 or \$100/ac/yr scenarios for both species, (scenario #4 – 7, Table 7 and 8).

Literature Cited

Bailey, R.L.; Zhao, B. 1998. GaPPS 4.20 Model. Warnell School of Forest Resources-UGA, Athens, GA.

Blevins, D.; Allen, H.L.; Colbert, S; Gardner, W. 1996. Nutrition management for longleaf pinestraw - Woodland Owner Notes. NC Coop. Ext Serv. Paper.

Clark III, A., Daniels, R.F. 2002. Wood quality of slash pine and its effect on lumber, paper, and other products. Presentation at the Slash Pine Symposium. Jekyll Island, GA. April 23-25, 2002.

Dickens, E.D. 1999. The effect of inorganic and organic fertilization on longleaf tree growth and pine straw production. In: Proceedings of the 10th Biennial So. Silvi. Res. Conf., Shreveport, LA, Feb 16-18, 1999. pp. 464-468.

Dickens, E.D. 2001. The effect of one-time biosolids application in an old-field loblolly pine plantation on diameter distributions, volume per acre, and value per acre. In: Proceedings of the 11th Biennial So. Silvi. Res. Conf., Knoxville, TN. March 19-22, 2001. pp. 15-19.

Doherty, B.A.; Teasley, R.J.; McKissick, J.C.; Givan, B. 2000. Nineteen ninety-nine farmgate value report. UGA CAES Center for Agribusiness and Econ. Dev., Center Staff Report No. 6. Athens, GA. 160 p.

Dubois, M.R.; McNabb, K.; Straka, T.K. 1999. Costs and cost trends for forestry practices in the South. *Forest Landowner Magazine*. March/April 1998. pp. 3-8.

Jokela, E.J.; Stearns-Smith, S.C. 1993. Fertilization of established southern pine stands: Effects of single and split nitrogen treatments. *SJAF* 17(3):135-138.

Martin, S.W.; Bailey, R.L.; Jokela, E.J. 1999. Growth and yield predictions for lower coastal plain slash pine plantations fertilized at mid-rotation. SJAF 23(1): 39-45.

Morris, L.A.; Jokela, E.J.; O'Connor, J.B., Jr. 1992. Silvicultural guidelines for pinestraw management in the SE US. GA Forest Res. Paper #88. GFC, Macon, GA. 11 p.

NCSFNC. 1998. North Carolina State University Forest Nutrition Coop - 26th Annual report. 23 p. School of Forest Resources, NCSU, Raleigh, NC.

Pienaar L.V.; Rheney, J.W. 1996. Potential productivity of intensively managed pine plantations - Final Report. The GA Consortium for Tech. Competitiveness in Pulp and Paper. 41 p.

Peinaar, L.V.; Shiver, B.D.; Rheney, J. W. 1996. Yield prediction for mechanically site-prepared slash pine plantations in the Southeastern Coastal Plain. PMRC Tech. Rep. 1996-3. 57 p.

Shiver, B.D.; Rheney, J.W.; Hitch, K.L.. 1999. Loblolly pine outperforms slash pine in southeast Georgia and northern Florida. SJAF 24(1) pp. 31-36.

TM-S 2004. Timber Mart So stumpage prices – 1st quarter South Georgia 2004. UGA-WSFR, Athens, GA 30602-2152.

Table 1. Product class specifications.

Product/Item	Pulpwood	Chip-N-Saw	Sawtimber
Small end diameter (inches)	3	6	10
Minimum length (feet)	5	8	8
Length Increment (feet)	1	4	8

Table 2. Product prices, cash and net (net of property taxes and marketing costs) per cord stumpage prices used in the profitability analysis of slash and loblolly scenarios, Georgia State average, price per ton (1stQ TM-S 2004).

Item, Price level	Cash or net	Pulpwood (\$/Ton)	Chip-N-Saw (\$/Ton)	Sawtimber (\$/Ton)
Low	cash	5.04	21.36	35.91
	net	4.51	19.12	32.14
Medium	cash	6.42	25.80	40.97
	net	5.75	23.09	36.51

Table 3. Fertilizer costs, \$100/acre/periodic application, per acre cost levels expressed as present values and annual equivalent values (AEV), as used in the profitability analysis for the 33-year slash and loblolly scenarios discounted at 8%.

Rotation (yrs)	Applied (yrs)	Present value of a periodic cost (\$/ac)	Annual equivalent value of the periodic cost (\$/ac/yr)
33	6	63.02	5.47
	6, 16	92.21	8.01
	6, 25	77.62	6.74
	16, 25	43.79	3.80
	6, 16, 25	106.81	9.28
	25	14.60	1.27

Table 4. Pine straw periodic per acre income levels expressed as present values and annual equivalent values (AEV) as used in the profitability analysis of slash and loblolly pine scenarios over the 33-year rotation discounted at 8%.

Item Rotation	Thin Scenario	Periodic income/ac/yr. raked (\$/ac)	Present value of periodic income (\$/ac)	AEV of periodic income (\$/ac/yr)
33 yrs	Thin at age 15 & 25 years	50 & 0 ¹	140.64	12.21
		100 & 0 ¹	281.28	24.43
		100 & 50 ²	351.64	30.54

¹ With thinning, pinestraw raked in years 8-14.

² With thinning, pinestraw raked in years 8-14 and 17-23.

Table 5. Clean-up cost, in year 16, after thinning of slash and loblolly pine scenarios over the 33-year rotation, expressed as present values and annual equivalent values (AEV) as used in the profitability analysis discounted at 8%.

Rotation	Clean-up cost in year 16 (\$/ac)	Present value of clean-up cost (\$/ac)	AEV of clean-up cost (\$/ac/yr)
33 years	70	20.43	1.77

Table 6. Stand management cost including active stand management, fire protection, and prescribed fire for loblolly and slash plantations 33-year rotation discounted at 8%.

Rotation	Management cost (\$/ac/yr)	Present value of Management cost year 1, (\$/ac)
33 years	9	103.62

Table7. A comparison of **slash pine plantation** management scenarios¹ under a **33-year rotation** with thinning in years 15 and 24 and their effect on economic variables, with site prep and plant (SP&PL) cost of **\$125, \$250, and \$375/acre.**

Treatments						SP&PL @ \$125		SP&PL @ \$250		SP&PL @ \$375	
Scenario # Fert. @ yr.	Pine straw (\$/ac)	% PW	% CNS	% St	MAI ² Ton, Cord	NR ³ (\$/ac)	ROR ⁴ (%)	NR ³ (\$/ac)	ROR ⁴ (%)	NR ³ (\$/ac)	ROR ⁴ (%)
1 N		35	35	30	5.26, 1.91	3100	9.86	2975	7.91	2850	6.69
2 Y, 25	N	34	34	32	5.58, 2.02	3267	10.06	3142	8.13	3017	6.92
3 Y, 16, 25		33	32	35	5.86, 2.13	3465	9.91	3340	8.11	3215	6.96
4 N	50 & 0 ⁵	34	35	30	5.26, 1.91	3450	12.88	3325	9.75	3200	8.08
5 Y, 6	100 & 0 ⁵	34	34	32	5.58, 2.02	3967	15.52	3842	11.28	3717	9.26
6 Y, 6, 25		33	32	35	5.86, 2.13	4165	15.32	4040	11.30	3915	9.34
7 Y, 6, 16, & 25	100 & 50 ⁶	32	31	37	6.26, 2.27	4736	16.89	4611	12.22	4486	10.06

Table 8. A comparison of **loblolly pine plantation** management scenarios¹ under a **33-year rotation** with thinning in years 15 and 24 and their effect on economic variables, with site prep and plant (SP&PL) cost of **\$125, \$250, and \$375/acre.**

Treatments						SP&PL @ \$125		SP&PL @ \$250		SP&PL @ \$375	
Scenario # Fert. @ yr.	Pine straw (\$/ac)	% PW	% CNS	% St	MAI ² Ton, Cord	NR ³ (\$/ac)	ROR ⁴ (%)	NR ³ (\$/ac)	ROR ⁴ (%)	NR ³ (\$/ac)	ROR ⁴ (%)
1 N		34	36	29	5.95, 2.15	3529	10.41	3404	8.42	3279	7.18
2 Y, 25	N	34	34	32	6.45, 2.34	3886	10.47	3761	8.57	3636	7.38
3 Y, 16, 25		33	32	35	6.99, 2.54	4237	10.94	4112	9.02	3987	7.80
4 N	50 & 0 ⁵	34	36	29	5.95, 2.15	3879	13.35	3754	10.20	3629	8.52
5 Y, 6	100 & 0 ⁵	34	34	32	6.45, 2.34	4586	15.66	4461	11.56	4336	9.59
6 Y, 6, 25		33	32	35	6.99, 2.54	4937	16.26	4812	12.12	4687	10.10
7 Y, 6, 16, & 25	100 & 50 ⁶	32	31	37	7.38, 2.68	5581	17.31	5456	12.81	5331	10.67

¹ Uninflated, 8% discount rate, before taxes, GaPPS v 4.20.

² MAI = Mean Annual Increment of wood growth for 33 yrs., tons & cords/A/yr.

³ NR = Net Revenue, revenues – costs (in today's \$s).

⁴ ROR = Rate of Return (percent).

⁵ With thinning, pinestraw raked years 8-14.

⁶ With thinning, pinestraw raked years 8-14 and 17-23.





Figure 1. Slash and loblolly pine 33-year rotation wood flow for scenario #1 and #4. (#1) thin @ age 15- and 24-years to 65 ft²/ac, no straw, no fertilization; (#4) thin @ age 15- and 24-years to 65 ft²/ac, rake pine straw from age 8- through age 14-years, no fertilization, (PW=pulpwood; CNS=chip-n-saw; ST=sawtimber; MAI=mean annual increment).





Figure 2. Slash and loblolly pine 33-year rotation wood flow for scenario #2 and #5. (#2) thin @ age 15- and 24-years to 65 ft²/ac, no straw, fertilize @ age 25-years; (#5) thin @ age 15- and 24-years to 65 ft²/ac, pine straw @ \$100/ac/yr from age 8- through age 14-years, fertilize @ age 6-years, (PW=pulpwood; CNS=chip-n-saw; ST=sawtimber; MAI=mean annual increment).





Figure 3. Slash and loblolly pine 33-year rotation wood flow for scenario #3 and #6. (#3) thin @ age 15- and 24-years to 65 ft²/ac, no straw, fertilize @ ages 16- and 25-years; (#6) thin @ age 15- and 24-years to 65 ft²/ac, pine straw @ \$100/ac/yr from age 8- through age 14-years, fertilize @ age 6- and 25-years (PW=pulpwood; CNS=chip-n-saw; ST=sawtimber; MAI=mean annual increment).