

Herbaceous Weed Control Recommendations for Planted Longleaf Sites

Revised 9 March 2012

Pat Minogue – Silviculture Assistant Professor – University of Florida School of Forest Resources and Conservation, David J. Moorhead – Silviculture Professor, and E. David Dickens – Forest Productivity Professor, The University of Georgia - Warnell School of Forestry and Natural Resources

- ◆ Herbaceous weed control (HWC) during the spring following planting longleaf can be critical to growth and survival on old-field, pasture, hayfield, and cut-over sites, particularly in droughty years.
- ◆ HWC can increase percentage of seedlings out of the grass stage by the 2nd year.
- ◆ A four to six foot wide band is often as effective as a broadcast treatment.
- ◆ Wait at least 2 months and with 1 to 2 inches of new feeder root growth from 2 to 3 or more lateral roots to spray herbicides over-the-top of longleaf after planting.

HERBICIDES FOR CONTROL OF BROADLEAF WEEDS AND GRASSES

(all treatments applied over-the-top of pines unless noted otherwise)

ARSENAL® (BASF; 53% imazapyr; 4 lb per gal)

- ◆ Very effective on perennial grasses, including difficult to control species like
- ◆ Bermudagrass, seedling Johnsongrass and Panicums
- ◆ Weak on broadleaf weeds in the composite group (see Oust XP®) and legumes
- ◆ Effective on established weeds
- ◆ Apply 4 – 6 fluid oz product per acre
- ◆ Do not add surfactant
- ◆ Optimum timing: Early post to post emergence of weeds (April - May)
- ◆ Grass and broadleaf weed control including, but not limited to: bahiagrass, barnyardgrass, bluegrass (annual, Kentucky), Bermudagrass, crabgrass, fescue, foxtail, Italian ryegrass, Johnsongrass, lovegrass, panicum (fall), sandbur, smooth brome, wild barley, wild oats, witchgrass, camphorweed, carpetweed, chickweed, clovers, cocklebur, dandelion, dogfennel, horseweed, goldenrod, knotweed, lambsquarters, milkweed, ragweed (common, giant), pepperweed, pigweed, plantain, pokeweed, purslane, pusley (Florida), shepard's purse, stinging nettle, sowthistle, annual spurge, sunflower, tansymustard, wild carrot, wild parsnip, wild turnip

OUST® XP® (DuPont; 75 % sulfometuron methyl)

- ◆ Very effect on a broad spectrum of broadleaf weeds, including composites

- ◆ Weak on perennial grass including Bermudagrass, broomsedge, and some Panicums (see Arsenal AC®)
- ◆ 1st Year plantings: apply 2-4 oz Oust product per acre
- ◆ Optimum timing: Pre-emergence (March-April)
- ◆ Do not use Oust when soil pH > 6.2
- ◆ Grass & broadleaf control including, but not limited to: chickweed, crabgrass, dogfennel, fescue, fireweed (willowweed), goldenrod, horseweed, Kentucky bluegrass, nutsedge (yellow), Panicum (broadleaf), pokeweed, ragweed, shepherd's purse, white snakeroot, yellow sweetclover, annual bluegrass, barnyardgrass, foxtail barley, foxtail fescue, Italian ryegrass, jointed goatgrass, bromes (red, ripgut), reed canarygrass, signalgrass, yellow foxtail, mustard, pepperweed, pigweed, sunflower, vetch, wild carrot, wild oats

OUST XP (DuPont; 75% sulfometuron methyl) + **VELPAR L®** (DuPont; 25% hexazinone)

- ◆ Broad spectrum weed control of broadleaf weeds and most grasses, weak on Bermudagrass and broomsedge
- ◆ Hexazinone may cause pine seedling mortality on sandy sites and on sites with low organic matter (old-field sites), ensure proper calibration and follow label directions regarding appropriate rates for various soil types
- ◆ Do not use Oust when soil pH > 6.2
- ◆ Apply 2-4 oz Oust product + VELPAR L 2 - 3 pints (or Velpar DF 10 – 16 oz product) per acre depending on soil texture (see product label)
- ◆ Optimum timing: Pre to early post emergence of weeds (March - early May)
- ◆ Use low rate of Oust + Velpar L or Velpar DF on coarse textured (sand, loamy sand, and sandy loam) soils and where soils are low in organic matter (see label)
- ◆ Grass & broadleaf control including, but not limited to: chickweed, crabgrass, dogfennel, fescue, fireweed (willowweed), goldenrod, horseweed, Kentucky bluegrass, nutsedge (yellow), Panicum (broadleaf), pokeweed, ragweed, shepherd's purse, white snakeroot, yellow sweetclover, annual bluegrass, barnyardgrass, foxtail barley, foxtail fescue, Italian ryegrass, jointed goatgrass, bromes (red, ripgut), reed canarygrass, signalgrass, yellow foxtail, mustard, pepperweed, pigweed, sunflower, vetch, wild carrot, wild oats, asters, brackenfern, fleabane

OUSTAR® (DuPont; 11.8% sulfometuron methyl and 63.2% hexazinone)

- ◆ Similar to Oust XP + Velpar products as above, but in a packaged mixture
- ◆ The ratio of active ingredients is set; hexazinone rate tend to be too high on sandy and low soil organic matter sites
- ◆ 1st Year weed control application product rates per acre:
 - 10-12 oz Course textured soils (sand, loamy sand, sandy loam)
 - 12-16 oz Medium textured soils (loam, sandy clay loam, silt loam)
 - 16-19 oz Fine textured soils (clay loam, sandy clay, silty clay loam, silty clay)
- ◆ After 1st year weed control application product rates per acre:
 - 12-16 oz Course textured soils

16-19 oz Medium textured soils

18-24 oz Fine textured soils

- ◆ Do not use Oustar when soil pH > 6.2
- ◆ Optimum timing: Pre to early post emergence (March - early May)
- ◆ Grass & broadleaf control including, but not limited to: chickweed, crabgrass, dogfennel, fescue, fireweed (willowweed), goldenrod, horseweed, Kentucky bluegrass, nutsedge (yellow), Panicum (broadleaf), pokeweed, ragweed, shepherd's purse, white snakeroot, yellow sweetclover

VELPAR DF (DuPont; 75% hexazinone)

- ◆ May cause mortality where excessive rates are applied on sandy soils and/or soils with low organic matter, ensure proper sprayer calibration to apply precise herbicide rate per acre, following label recommendations regarding specific herbicide rates for various soil types
- ◆ 1st Year weed control application product rates per acre (the same amounts can be applied in years 2, 3, and 4):
 - 1 1/3 lb Course textured soils (loamy sand, sandy loam)
 - 1 1/3 – 1 1/2 lb Medium textured soils (loam, sandy clay loam, silt, silt loam)
 - 1 1/2 - 1 4/5 lb Fine textured soils (sandy clay, silty clay loam, silty clay, clay, clay loam)
- ◆ Weed control application product rates per acre for established trees (≥ 4-yrs-old):
 - 1 1/3 – 1 2/3 lb Course textured soils
 - 1 2/3 – 2 1/3 lb Medium textured soils
 - 2 1/3 – 2 2/3 lb Fine textured soils
- ◆ Optimum timing: Pre to early post emergence (March – early May)
- ◆ Grass & broadleaf control including, but not limited to: Asters, barnyardgrass, annual bluegrass, brackenfern, bromegrass, fleabane, foxtail, horseweed, ragweed, ryegrass

VELPAR L (DuPont; 25% Hexazinone)

- ◆ May cause mortality where excessive rates are applied on sandy soils and/or soils with low organic matter, ensure proper sprayer calibration to apply precise herbicide rate per acre, following label recommendations regarding specific herbicide rates for various soil types
- ◆ 1st Year weed control application product rates per acre (the same amounts can be applied in years 2, 3, and 4):
 - 21 to 32 oz Course textured soils (loamy sand, sandy loam)
 - 24 to 40 oz Medium textured soils (loam, sandy clay loam, silt, silt loam)
 - 28 to 48 oz Fine textured soils (clay, clay loam, sandy clay, silty clay loam, silty clay)
- ◆ After 4th year weed control application product rates per acre:
 - 21 to 40 oz Course textured soils
 - 28 to 56 oz Medium textured soils
 - 36 to 64 oz Fine textured soils

- ◆ Optimum timing: Pre to early post emergence (March - early May)
- ◆ Grass & broadleaf control including, but not limited to: Asters, barnyardgrass, annual bluegrass, brackenfern, bromegrass, fleabane, foxtail, horseweed, ragweed ryegrass

Also consider:

(1) The herbicide purchaser and applicator need to look closely at the herbicide label to make sure that the herbicide product used is: (a) labeled for the particular use site (e.g. “for use in forest sites”, “for use in conifer plantations”, etc.) (b) labeled for the pine crop species (or genus in some cases), and (c) labeled for the particular application (“herbaceous weed control in pine plantations”, “herbaceous release”, etc.). Herbicide products, even those with the same trade name (such as “Roundup”), may have different amounts of the active ingredient per gallon, so always follow the rates specified on the label of the particular container being used.

(2) Remember rates are per acre treated. Here are two examples of calculating the herbicide needed for a banded herbicide application using 10 oz Oustar product per acre treated, assuming 12 feet between the rows: (a) spraying a 4 foot wide band, 10 oz Oustar will take care of 3 acres total land area. In effect, one acre is banded and two acres are untreated, so herbicide is applied to one-third of the area. A total of 400 oz (25 lbs) of Oustar would be needed for a 120 acre field where one-third the area is treated in bands. (b) When spraying a 6 foot wide band on rows 12 feet apart, 10 oz Oustar will take care of 2 acres total land area. One acre is banded and one acre is untreated, so herbicide is applied to half the area. A total of 600 oz (37.5 lbs) of Oustar would be needed for a 120 acre field where herbicide is applied to one-half the area in bands.

(3) There are generics for some of the above listed herbicides and various product labels for the same active ingredient do vary. Read the product labels to make sure that your intended use is consistent with labeling.

Please read and follow all label recommendations. Inclusion of a product trade name or a company name in this publication does not constitute an endorsement of a product or a company, as other products manufactured by different companies might be equally suited for the intended herbicide use.

HERBICIDES FOR CONTROL OF ANNUAL & PERENNIAL GRASSES ONLY

- (1) When restoring native grasses concurrently to establishing longleaf pine, be careful not to apply the following “grass herbicides” in the Warm Season Native Grass (WSNG) planting area after grasses have emerged.
- (2) All grass control herbicides listed below are postemergence, foliar active herbicides.
- (3) Best control for all grass species is obtained when grasses are in an early growth stage. For Texas panicum, apply when the grass is less than 4 inches tall. For Bermudagrass two applications are usually needed; the first when less than 6 inches tall and a second when re-growth is less than 4 inches. Multiple applications are also needed for Johnsongrass.
- (4) Herbicides in this group generally do not mix well with other herbicide products. However, it is very important to add surfactants (wetting agents) to improve plant uptake. See information below and product labels for details.
- (5) Herbicide spray solution (water) volumes are typically between 10 to 20 gallons per

- sprayed acre (GPA) with a range 5-40 GPA; read label for specifics.
- (6) Do not apply herbicides when pine seedlings and desirable grasses are under drought or other stress
 - (7) Do not apply herbicides when rainfall is expected within one hour.

ENVOY® PLUS (Valent; 12.6% clethodim, 0.97 lb clethodim per gallon, contains petroleum distillates)

- ◆ Apply 9 to 16 fluid oz per acre for annual grasses, 12 to 32 oz/acre for perennial grasses
- ◆ Add crop oil concentrate which contains at least 15% emulsifier at 1% volume/volume (1 qt per 25 gallons spray solution, but no less than 1 pint per acre) or non-ionic surfactant at 0.25% volume/volume (1 qt per 100 gallons)
- ◆ Apply in 10 to 40 gallons of water per acre

ENVOY® (Valent; 12.6% clethodim, 0.94 lbs clethodim per gallon, contains petroleum distillates)

- ◆ Apply 13 – 34 fluid oz product per acre,
- ◆ Use a lower dose for annual grasses, a higher dose for perennial grasses
- ◆ Add crop oil concentrate with at least 15% emulsifier at 1% volume/volume (1 qt per 25 gallons, but no less than 1 pint per acre) or non-ionic surfactant at 0.25% volume/volume (1 qt per 100 gallons)

FUSILADE® DX (Syngenta; 24.5% fluazifop-P-butyl, 2 lb per gallon fluazifop-P-butyl)

- ◆ Apply 16-24 fluid oz product per acre per application
- ◆ Use a lower dose for annual grasses, a higher dose for perennial grasses
- ◆ Add 1% crop oil concentrate (1 quart per 25 gal) or 0.25% nonionic surfactant (1 quart per 100 gal)
- ◆ Do not apply more than 72 fluid oz Fusilade DX per acre, per season

ARROW® 2EC (Makhteshim Agan of North America (MANA); 26.4% clethodim, 2.0 lbs clethodim per gallon, contains petroleum distillates)

- ◆ Apply 6 to 8 fluid oz product per acre for annual grasses and 8 to 16 oz/acre for perennial grasses
- ◆ Add crop oil concentrate which contains at least 15% emulsifier at 1% volume/volume (1 qt per 25 gallons spray solution, but no less than 1 pint per acre) or non-ionic surfactant at 0.25% volume/volume (1 qt per 100 gallons)

WEED TOLERANCE TO SELECTED HERBICIDES

Herbicide	Weeds
ARSENAL	sicklepod, tropic croton, blackberry, most legumes
ARROW 2EC, ENVOY, ENVOY Plus	All broadleaf weeds
FUSILADE DX	All broadleaf weeds
OUST	bermudagrass, croton, Johnsongrass, trumpetcreeper, broomsedge
VELPAR	bermudagrass, broomsedge, cocklebur, Johnsongrass, sicklepod, trumpetcreeper, morningglory

GRASS WEED RESPONSE TO HERBICIDES

WEED	^aFluazifop-P-butyl	^bClethodim
Perennial Grasses		
bermudagrass	G - E	G - E
Bahiagrass	G	
Johnsongrass (rhizome)	E	E
tall fescue	F	
nutsedge	P	P
Annual Grasses		
broadleaf signalgrass	E	E
crowfootgrass	F	G
crabgrass	F	G
fall panicum	G	G
goosegrass	F - G	F - G
Johnsongrass (seedling)	G - E	E
sandbur	G	G - E
Texas panicum	G - E	G

E=Excellent (>90% control); G=Good(80-89% control); F=Fair(70-79% control); P=Poor(<70%). Citations: Guillebeau, P. (ed). 2005 & 2008. Georgia Pest Mgmt. Handbook. Special Bulletin 28. UGA Coop. Ext. Service. Athens, GA 30602. pp. 112, 156-57, 198-99. ^aFluazifop-P-butyl rating based on Fusilade DX trials, ^bClethodim ratings based on Select® and Arrow® trials.

Old-field non-scalped post-plant herbaceous weed control timing considerations for the Georgia Coastal Plain and Central to Northern Florida

Soil drainage class	Pre- to early post emergence herbicide	Early post to post emergence herbicide
Somewhat excessively to excessively well	March	March to early April
Moderately well to well	March to early April	mid-March to mid-April
Poorly to somewhat poorly	April to early May	mid-April to mid-May

* Do not apply herbicides over longleaf for at least 2 months after planting and when there is 1 to 2 inches of new feeder root growth off 2 or more lateral roots and wait at least 1 to preferably 2 months after planting for slash pine.

** For scalped sites, herbicides applied from mid-April into mid- to late May have given good survival and growth results based on recent studies as long as seedlings are not under stress (especially drought stress).

Organization of GA (FL, AL, and SC in some cases) Coastal Plain Soil Series in Management Groups (Larry Morris "Forest soils and management decisions" workshop 2005)

Subsoil Type:		None	Loamy	Clayey	Spodic	
Drainage	Surface Depth (inches)	(Sandy to loamy sand)			+ Arg	- Arg.
Very poorly		Rutledge	Torhunta Surrency	Bayboro		Murville Wesconnet
Poorly to Somewhat Poorly	0-10	Chiple Osier Scranton	Rains	Bladen Coxville Grady	Rigdon	Ridgeland
	10-20		Lynchburg		Mascotte Sapelo	Leon Mandarin
	20-40		Pelham	Nanhunta		
	40-80		Albany Plummer	Kanapaha		Hurricane Pottsburg
Moderately Well to Well Drained	0-10	Resota Pactolus Ortega	Goldsboro	Faceville Nankin Greenville	Onslow	
	10-20		Tifton Dothan		Seagate	Baymeade
	20-40		Lucy Fuquay Stilson	[Dotted pattern]		Echaw
	40-80		Bonifay			
Somewhat to Excess. Well	40-80	Lakeland Kershaw	Troup			Rimini Kureb

Citation: Minogue, P. D.J. Moorhead, and E.D. Dickens. 2012 (Revised). Herbaceous weed control for planted longleaf sites. www.forestproductivity.net 7 p.

Athens, Georgia 30602-2152

Phone: 706.542.6819 • fax: 706.542.5073

An Equal Opportunity/Affirmative Action Institution

In compliance with federal law, including the provisions of Title IX of the Education Amendments of 1972, Title VI of the Civil Rights Act of 1964, Sections 503 and 504 of the Rehabilitation Act of 1973, and the Americans with Disabilities Act of 1990, the University of Georgia does not discriminate on the basis of race, sex, religion, color, national or ethnic origin, age, disability, or military service in its administration of educational policies, programs, or activities; its admissions policies; scholarship and loan programs; athletic or other University-administered programs; or employment. In addition, the University does not discriminate on the basis of sexual orientation consistent with the University non-discrimination policy. Inquiries or complaints should be directed to the director of the Equal Opportunity Office, Peabody Hall, 290 South Jackson Street, University of Georgia, Athens, GA 30602 Telephone 706-542-7912 (V/TDD). Fax 706-542-2822.